

EXHIBIT A

Behavioral Health Care > Care Guidelines for Behavioral Health > Posttraumatic Stress Disorder > Posttraumatic Stress Disorder - Annotated Bibliography > Posttraumatic Stress Disorder: Residential Care

Posttraumatic Stress Disorder: Residential Care

ORG: B-013-RES (BHG)

[Link to Codes](#)

MCGTM
Behavioral Health Care
17th Edition

- Residential Care Admission and Alternatives
 - Clinical Indications for Admission to Residential Care
 - Alternatives to Admission
 - Alternative Care Planning
- Residential Care Treatment
 - Recovery Course
 - Residential Care Planning
 - Discharge Criteria
- Discharge
 - Discharge Planning
 - Discharge Destination
 - Usual
 - Alternate
- Annotated Bibliography
- References
- Footnotes
- Codes

Residential Care Admission and Alternatives

[Return to top of Posttraumatic Stress Disorder: Residential Care - BHG](#)

Clinical Indications for Admission to Residential Care

[Return to top of Posttraumatic Stress Disorder: Residential Care - BHG](#)

[[Expand All](#) / [Collapse All](#)]





- Admission is indicated because of **1 or more** of the following:
 - ☐ Admission to Residential Acute Level of Care is judged appropriate as indicated by **ALL** of the following (A):
 - Around-the-clock behavioral care is necessary for treatment because of **1 or more** of the following:
 - Imminent danger to self because of **1 or more** of the following (1)(4)(5)(6):
 - Imminent risk for recurrence of Suicide attempt or act of serious self Harm as indicated by **ALL** of the following:
 - Very recent Suicide attempt or deliberate act of serious self Harm
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for suicide or serious self Harm
 - Command auditory hallucinations for suicide or serious self Harm
 - Persistent Thoughts of suicide or serious Harm to self, or suicide trigger state without formed thoughts, that cannot be adequately monitored at lower level of care because of **1 or more** of the following (A)(7)(8):
 - Insufficient behavioral care (eg, facility, provider) availability
 - Inadequate patient support system
 - Patient characteristics such as high impulsivity, unreliability, or extreme agitation with desperation
 - Ruminative flooding; uncontrollable and overwhelming profusion of negative thoughts
 - Frantic hopelessness; fatalistic conviction that life will not improve along with oppressive sense of entrapment and doom
 - Imminent danger to others due to **1 or more** of the following (5)(9)(10)(11):
 - Imminent risk for recurrence of an attempt to seriously Harm another as indicated by **ALL** of the following:
 - Very recent attempt to seriously Harm another
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for homicide or serious Harm to another
 - Command auditory hallucinations or paranoid delusions contributing to risk for homicide or serious Harm to another
 - Persistent thoughts of homicide or serious Harm to another that cannot be adequately monitored at lower level of care because of **1 or more** of the following (A):
 - Insufficient behavioral care (eg, facility, provider) availability
 - Inadequate patient support system
 - Patient characteristics such as high impulsivity or unreliability
 - Life-threatening inability to perform self-care activity (eg, self neglect with inability to provide for self at lower level of care) (9)(10)
 - Severe disability or disorder requiring acute residential intervention as indicated by **ALL** of the following:

- Severe behavioral health disorder-related symptoms or condition are present as indicated by **1 or more** of the following (4)(5)(12)(13)(14):
 - Major dysfunction in daily living (eg, social, interpersonal, occupational functioning)
 - Severe problem with cognition, memory, or judgment (15)
 - Severe symptoms (eg, hallucinations, delusions, other acute psychotic symptoms, mania, extreme agitation or anxiety) (16)(17)
 - Patient management at highest nonresidential level of care has failed or is not feasible until acute intervention or modification is initiated.
 - Severe comorbid substance use disorder that must be controlled (eg, abstinence necessary) to achieve stabilization of primary psychiatric disorder (9)(18)(19)(20)
 - Patient has currently stabilized during inpatient treatment stay for severe symptoms or behavior and requires structured setting with continued around-the-clock behavioral care.
 - No exclusions to treatment; situation and expectations are appropriate for level as indicated by **ALL** of the following (1)(2)(3)(5)(9)(13)(4)(14):
 - Recommended treatment is necessary, appropriate, and not feasible at a lower level of care (ie, documented behavior, symptoms, or risk judged not appropriate for partial hospital, IOP, or acute outpatient care).
 - Very short-term crisis intervention and resource planning for further care at a nonresidential level is unavailable or judged inappropriate.
 - Patient is willing to participate in treatment within a highly structured setting voluntarily.
 - No anticipated need for physical restraint, seclusion, or other involuntary control (eg, patient not actively violent)
 - No need for around-the-clock medical or nursing care
 - Patient has sufficient cognitive capacity to respond to planned individual and group treatment components.
 - Adequate response (eg, stabilization for nonresidential level of care) to planned treatment is expected within a limited time period.
- ☐ Admission to Residential Acute Level of Care for Child or Adolescent is judged appropriate as indicated by **ALL** of the following (4):
- Around-the-clock behavioral care is necessary for treatment because of **1 or more** of the following (21):
 - Imminent danger to self because of **1 or more** of the following (6)(22)(23):
 - Imminent risk for recurrence of Suicide attempt or act of serious self Harm as indicated by **ALL** of the following:
 - Very recent Suicide attempt or deliberate act of serious self Harm
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for suicide or serious self Harm
 - Command auditory hallucinations for suicide or serious self Harm
 - Dangerous behavior risk, persistent Thoughts of suicide or serious Harm to self, or suicide trigger state without formed thoughts, that cannot be adequately monitored at lower level of care because of **1 or more** of the following (8)(22):
 - Insufficient child or adolescent behavioral care (eg, facility, provider) availability
 - Severe conflict in family environment or other inadequacy in patient support system
 - Patient characteristics such as high impulsivity, unreliability, or extreme agitation with desperation
 - Ruminative flooding; uncontrollable and overwhelming profusion of negative thoughts
 - Frantic hopelessness; fatalistic conviction that life will not improve along with oppressive sense of entrapment and doom
 - Imminent danger to others due to **1 or more** of the following (5)(10)(11)(24):
 - Imminent risk for recurrence of an attempt to seriously Harm another as indicated by **ALL** of the following:
 - Very recent attempt to seriously Harm another
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for homicide or serious Harm to another
 - Command auditory hallucinations or paranoid delusions contributing to risk for homicide or serious Harm to another
 - Persistent thoughts of, or violent impulsive acts that could likely result in, homicide or serious Harm to another that cannot be adequately monitored at lower level of care because of **1 or more** of the following (8):
 - Insufficient child or adolescent behavioral care (eg, facility, provider) availability
 - Severe conflict in family environment or other inadequacy in patient support system
 - Patient characteristics such as high impulsivity or unreliability
 - Life-threatening inability to receive adequate care from caretakers (5)(10)
 - Severe disability or disorder requiring acute residential intervention as indicated by **ALL** of the following:
 - Severe behavioral health disorder-related symptoms or condition are present as indicated by **1 or more** of the following (2)(3)(5)(12)(13)(14):
 - Major dysfunction in daily living (eg, family, interpersonal, school functioning)
 - Severe problem with cognition, memory, or judgment
 - Severe symptoms (eg, hallucinations, delusions, other acute psychotic symptoms, mania, severe autistic behaviors) (16)(17)(25)
 - Severe behavior risk (affective dysregulation) characteristics indicated by **1 or more** of the following:
 - Evidence of severely diminished ability to assess consequences of own actions (eg, acts of severe property damage)
 - Frequent extreme external (extreme angry outbursts) or internal (extreme sulking and rumination) anger manifestations
 - High levels of family conflict
 - Patient management at highest nonresidential level of care has failed or is not feasible until acute intervention or modification is initiated.

- Severe comorbid substance use disorder that must be controlled (eg, abstinence necessary) to achieve stabilization of primary psychiatric disorder (19)(20)
- Patient has currently stabilized during inpatient treatment stay for severe symptoms or behavior and requires structured setting with continued around-the-clock behavioral care.
- No exclusions to treatment: situation and expectations are appropriate for level as indicated by ALL of the following (1)(2)(3)(5)(9)(13)(21)(22):
 - Recommended treatment is necessary, appropriate, and not feasible at a lower level of care (ie, documented behavior, symptoms, or risk judged not appropriate for partial hospital, IOP, or acute outpatient care).
 - Very short-term crisis intervention and resource planning for further care at a nonresidential level is unavailable or judged inappropriate.
 - Patient has, at the least, minimal motivation to participate in treatment within a highly structured setting at the direction of parent or guardian.
 - No anticipated need for physical restraint, seclusion, or other involuntary control (eg, patient not actively violent)
 - No need for around-the-clock medical or nursing care
 - Patient has sufficient cognitive capacity to respond to planned individual and group treatment components. (14)
 - Adequate response (eg, stabilization for nonresidential level of care) to planned treatment is expected within a limited time period.

Alternatives to Admission

Return to top of *Posttraumatic Stress Disorder: Residential Care - BHG*

- Alternatives include (4)(7)(26)(27):
 - Acute outpatient care: appropriate if around-the-clock behavioral care is not necessary, and needed type and frequency of treatment are available in office or clinic setting. (28)(29)(30)(31) See *Posttraumatic Stress Disorder: Acute Outpatient Care* .
 - Intensive outpatient program: appropriate if around-the-clock behavioral care is not necessary, and needed type and frequency of treatment are available in intensive outpatient program but not in office or clinic setting. (32)(33) See *Posttraumatic Stress Disorder: Intensive Outpatient Program* .
 - Partial hospital program: appropriate if around-the-clock behavioral care is not necessary, and support is available to provide any needed monitoring of patient's condition when partial hospital program is closed. (32)(34)(35) See *Posttraumatic Stress Disorder: Partial Hospital Program* .
 - Care for higher-risk or more severely ill patient in inpatient care. See *Posttraumatic Stress Disorder: Inpatient Care* .

Alternative Care Planning

Return to top of *Posttraumatic Stress Disorder: Residential Care - BHG*

- Care planning needs for patient not requiring residential admission may include:
 - Prompt visit to a psychiatrist for assessment and possible pharmacotherapy (29)(36)(37)(38)(39)(40)
 - Referral for trauma-focused individual psychotherapy (29)(42)(43)(44)(45)(46)(47)
 - Medical care visit (eg, primary care) to evaluate comorbidity or clear for pharmacotherapy
 - Discharge planning as appropriate. See *Discharge Planning* in this guideline.

Residential Care Treatment

Return to top of *Posttraumatic Stress Disorder: Residential Care - BHG*

Recovery Course

Return to top of *Posttraumatic Stress Disorder: Residential Care - BHG*

Stage	Clinical Status	Interventions	Medications	Evaluation	Therapy
1	<ul style="list-style-type: none"> • Clinical indications met (E) • Begin discharge planning 	<ul style="list-style-type: none"> • Possible frequent safety checks (F) 	<ul style="list-style-type: none"> • Possible antidepressant (C) 	<ul style="list-style-type: none"> • Exploration of admission precipitants (G) • Psychiatric, social, medical, substance use, and traumatic events histories (H) • Mental status and physical examinations • Symptoms assessed multiple times per shift 	<ul style="list-style-type: none"> • Psychosocial interventions emphasizing admission precipitants (I) (daily)

2

- Possible frequent safety checks
- Medication review if prescribed
- **Evaluation completed and reviewed**
- Symptoms assessed multiple times per shift
- Psychosocial interventions emphasizing admission precipitants (daily)
- Clinical management and psychoeducation if medication is prescribed [4] (every other day)
- Parental component of psychosocial interventions if patient is child or adolescent (every third day)

3

- **No plan [M] for suicide or serious self-harm [L] for at least 24 hours**
- No need for frequent safety checks
- Medication review if prescribed
- Transition to patient check-in with staff
- Psychosocial interventions emphasizing barriers to discharge [I] (daily)
- Clinical management and psychoeducation if medication is prescribed (every other day)
- Parental component of psychosocial interventions if patient is child or adolescent (every third day)

4

- Risk status acceptable
- Functional status acceptable
- Complete discharge planning
- Discharge
- Review follow-up treatment and crisis plan with patient and supports [M]
- Medical needs manageable

(9)(28)(29)(30)(32)(36)(37)(39)(50)(51)

Recovery Milestones are indicated in **bold**.**Residential Care Planning**

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

- Residential evaluation and care needs may include:
 - Diagnostic test scheduling and completion, including [M] (49):
 - Urine drug screen [M]
 - Pregnancy test
 - Treatment and procedure scheduling and completion, including:
 - Psychosocial interventions emphasizing admission precipitants and barriers to discharge [I]
 - Clinical management and psychoeducation if medication is prescribed [4]
 - Parental component of psychosocial interventions if patient is child or adolescent(29)
 - Trauma-focused individual psychotherapy [D] [C]
 - Antidepressant medication [C]
 - Consultation, assessment, and other services scheduling and completion, including:
 - Substance abuse assessment(28)
 - Social services consultation for placement or housing
 - Conference with school for child or adolescent
 - Monitoring patient's status for deterioration and comorbid conditions; key items include:
 - Psychiatric comorbidity [M]
 - Substance withdrawal
 - Pre-existing medical comorbidity

Discharge Criteria

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

- Continued residential care is generally needed until 1 or more of the following(9):
 - Residential care no longer necessary due to adequate patient stabilization or improvement as indicated by **ALL** of the following(4)(7)(26) (27)(28):


- Risk status acceptable as indicated by ALL of the following:
 - Patient has not recently made a Suicide attempt or act of serious self Harm , or has had Sufficient relief of precipitants of any such action.
 - Absence of Current plan for suicide or serious self Harm for at least 24 hours
 - Thoughts of suicide , homicide, or serious Harm to self or to another are absent or manageable at available lower level of care.
 - Patient and supports understand followup treatment and crisis plan.
 - Provider and supports are sufficiently available at lower level of care.
 - Patient can participate (eg, verify absence of plan for harm) in needed monitoring.
- Functional status acceptable as indicated by 1 or more of the following:
 - No essential function FI is significantly impaired.
 - An essential function is impaired, but impairment is manageable at available lower level of care.
- Medical needs manageable as indicated by ALL of the following:
 - Adverse medication effects absent or manageable at available lower level of care
 - Medical comorbidity absent or manageable at available lower level of care
 - Substance withdrawal absent or manageable at available lower level of care
- Residential care no longer appropriate due to patient progress record or consent as indicated by 1 or more of the following:
 - Patient deterioration requires higher level of care.
 - Patient or guardian no longer consents to treatment.

Discharge

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

Discharge Planning

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

- Discharge planning needs may include(51):
 - Rapid decision and planning regarding next level of care; considerations include:
 - Plan for monitoring for dangerous ideation or behavior if necessary(7)(26)
 - Plan for assisting patient with self-care if necessary
 - Follow-up plan development with input from multiple providers, patient, and patient's supports(28)
 - Preparation of patient and supports for transition to lower level of care, including:
 - Ensure sufficient knowledge of(28)(29):
 - Patient's illness
 - Medication
 - Risk factors for relapse
 - Warning signs of relapse
 - Assess ability to work, attend school, and participate in usual activities.
 - Review crisis plan with patient and supports. (M)
 - Patient and caregiver education complete. See Posttraumatic Stress Disorder: Patient Education for Clinicians  SR.
 - Follow-up appointments, including:
 - Psychiatrist for pharmacotherapy and clinical management (C)
 - Therapist for trauma-focused psychotherapy (C)
 - Medical care visit (eg, primary care)
 - Referrals for community assistance and support, including:
 - Self-help or support groups for patient, family, and caregivers(28)
 - Community services for housing, financial, or transportation needs
 - Discharge medications and supplies, including:
 - Antidepressant medication
 - Medications for comorbid medical conditions

Discharge Destination

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG



Usual

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

- Acute outpatient care. See Posttraumatic Stress Disorder: Acute Outpatient Care  BHG.

Alternate

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

- Intensive outpatient program: appropriate if around-the-clock behavioral care is not necessary, and needed type and frequency of treatment are available in intensive outpatient program but not in office or clinic setting. See Posttraumatic Stress Disorder: Intensive Outpatient Program  BHG.
- Partial hospital program: appropriate if around-the-clock behavioral care is not necessary, and support is available to provide any needed monitoring of patient's condition when partial hospital program is closed. See Posttraumatic Stress Disorder: Partial Hospital Program  BHG.

Annotated Bibliography

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

See Posttraumatic Stress Disorder - Annotated Bibliography  BHG for a discussion of key literature.

References

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

1. Lloyd-Evans B, Slade M, Jagielska O, Johnson S. Residential alternatives to acute psychiatric hospital admission: systematic review. *British Journal of Psychiatry* 2009;195(2):109-17. DOI: 10.1192/bjp.bp.108.058347. [Context Link 1, 2, 3, 4] View abstract...
2. Mee-Lee D, Shulman GD, Fishman M, Gastfriend DR, Griffith JH. ASAM Patient Placement Criteria for the Treatment of Substance-Related Disorders, Second Edition-Revised. 2nd rev ed. Chevy Chase, MD: American Society of Addiction Medicine, 2001. [Context Link 1, 2, 3, 4]
3. Kleber HD, et al. Practice guideline for the treatment of patients with substance use disorders. 2nd ed. [Internet] American Psychiatric Association. 2007 Apr Accessed at: <http://psychiatryonline.org/guidelines.aspx>. [created 2006 May; accessed 2012 Sep 20] DOI: 10.1176/appi.books.9780890423363.141077. [Context Link 1, 2, 3, 4]
4. Gelenberg AJ, et al. Treatment of patients with major depressive disorder. 3rd edition [Internet] American Psychiatric Association. 2010 Oct 1 Accessed at: <http://psychiatryonline.org/guidelines.aspx>. [accessed 2012 Aug 28] DOI: 10.1176/appi.books.9780890423363.654001. [Context Link 1, 2, 3, 4, 5]
5. Lichtenberg P. The residential care alternative for the acutely psychotic patient. *Psychiatric Quarterly* 2011;82(4):329-41. DOI: 10.1007/s11126-011-9176-0. [Context Link 1, 2, 3, 4, 5, 6, 7, 8] View abstract...
6. Diesvold T, Bakkejord T, Hansen V, Nivison M, Sorgaard KW. Suicidality related to first-time admissions to psychiatric hospital. *Social Psychiatry and Psychiatric Epidemiology* 2012;47(3):419-25. DOI: 10.1007/s00127-011-0343-2. [Context Link 1, 2] View abstract...
7. Sudak HS. Suicide. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:2717-31. [Context Link 1, 2, 3, 4, 5]
8. Yaseen Z, Katz C, Johnson MS, Eisenberg D, Cohen LJ, Galyenker II. Construct development: The Suicide Trigger Scale (STS-2), a measure of a hypothesized suicide trigger state. *BMC Psychiatry* 2010;10:110. DOI: 10.1186/1471-244X-10-110. [Context Link 1, 2] View abstract...
9. Hawthorne WB, et al. A randomized trial of short-term acute residential treatment for veterans. *Psychiatric Services* 2005;56(11):1379-86. DOI: 10.1176/appi.ps.56.11.1379. [Context Link 1, 2, 3, 4, 5, 6, 7] View abstract...
10. Baron DA, Dubin WR, Ning A. Other psychiatric emergencies. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:2732-45. [Context Link 1, 2, 3, 4]
11. Myklebust LH, Sorgaard K, Rotvold K, Wynn R. Factors of importance to involuntary admission. *Nordic Journal of Psychiatry* 2012;66(3):178-82. DOI: 10.3109/08039488.2011.611252. [Context Link 1, 2] View abstract...
12. Brooker C, Ricketts T, Bennett S, Lemme F. Admission decisions following contact with an emergency mental health assessment and intervention service. *Journal of Clinical Nursing* 2007;16(7):1313-22. DOI: 10.1111/j.1365-2702.2007.01302.x. [Context Link 1, 2] View abstract...
13. Stewart SE, et al. Long-term outcome following intensive Residential Treatment of Obsessive-Compulsive Disorder. *Journal of Psychiatric Research* 2009;43(13):1118-23. DOI: 10.1016/j.jpsychires.2009.03.012. [Context Link 1, 2, 3, 4] View abstract...
14. Boyer DE, Kane C. Program evaluation of a community crisis stabilization program. *Archives of Psychiatric Nursing* 2010;24(6):387-96. DOI: 10.1016/j.apnu.2010.03.002. [Context Link 1, 2, 3, 4] View abstract...
15. Caplan JP, Rabinowitz T. An approach to the patient with cognitive impairment: delirium and dementia. *Medical Clinics of North America* 2010;94(6):1103-16. ix. DOI: 10.1016/j.mcna.2010.08.004. [Context Link 1] View abstract...
16. Lewis S, Escalona PR, Keith SJ. Phenomenology of schizophrenia. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:1433-50. [Context Link 1, 2]
17. Fochtmann LJ, Mojtabai R, Bromet EJ. Other psychotic disorders. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:1605-28. [Context Link 1, 2]
18. Farnen KC, McElroy S. Treatment response of bipolar and unipolar alcoholics to an inpatient dual diagnosis program. *Journal of Affective Disorders* 2008;106(3):265-72. [Context Link 1]
19. Kuehn BM. Integrated care key for patients with both addiction and mental illness. *Journal of the American Medical Association* 2010;303(19):1905-7. DOI: 10.1001/jama.2010.597. [Context Link 1, 2]
20. Torrens M, Gilchrist G, Domingo-Salvany A, psyCoBarcelona Group. Psychiatric comorbidity in illicit drug users: substance-induced versus independent disorders. *Drug and Alcohol Dependence* 2011;113(2-3):147-56. DOI: 10.1016/j.drugalcdep.2010.07.013. [Context Link 1, 2] View abstract...
21. Kober D, Martin A. Inpatient psychiatric, partial hospital, and residential treatment for children and adolescents. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3766-71. [Context Link 1, 2]
22. Wagner KD, Brent DA. Depressive disorders and suicide. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3652-62. [Context Link 1, 2, 3]
23. Screening and treatment for major depressive disorder in children and adolescents: US Preventive Services Task Force Recommendation Statement. *Pediatrics* 2009;123(4):1223-8. DOI: 10.1542/peds.2008-2381. [Context Link 1] View abstract...
24. Meuleners LB, Lee AH, Hendrie D. Interpersonal violence hospitalisations for adolescents: a population-based study. *Journal of Paediatrics and Child Health* 2010;46(11):886-90. DOI: 10.1111/j.1440-1754.2010.01818.x. [Context Link 1] View abstract...
25. Volkmar FR, Klin A, Schultz RT, Stale MW. Pervasive developmental disorders. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3540-59. [Context Link 1]
26. Velligan DI, et al. Strategies for addressing adherence problems in patients with serious and persistent mental illness: recommendations from the expert consensus guidelines. *Journal of Psychiatric Practice* 2010;16(5):306-24. DOI: 10.1097/J1.pra.0b00386626.98862.s0. [Context Link 1, 2, 3] View abstract...
27. Sharfstein SS. Goals of inpatient treatment for psychiatric disorders. *Annual Review of Medicine* 2009;60:393-403. DOI: 10.1146/annurev.med.60.042607.080257. [Context Link 1, 2] View abstract...
28. Practice guideline for the treatment of patients with acute stress disorder and posttraumatic stress disorder. [Internet] American Psychiatric Association. 2004 Nov Accessed at: <http://psychiatryonline.org/guidelines.aspx>. [accessed 2012 Aug 27] DOI: 10.1176/appi.books.9780890423363.52257. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9]
29. National Collaborating Centre for Mental Health. Post-traumatic stress disorder: the management of PTSD in adults and children in primary and secondary care. NICE Clinical Guidance CG26 [Internet] Royal College of Psychiatrists and the British Psychological Society. 2005 Mar Accessed at: <http://guidance.nice.org.uk/>. [accessed 2012 Aug 27] [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]

30. De Bellis MD, Van Dillen T. Childhood post-traumatic stress disorder: an overview. *Child and Adolescent Psychiatric Clinics of North America* 2005;14(4):745-72. ix. DOI: 10.1016/j.chc.2005.05.006. [Context Link 1, 2, 3] View abstract...
31. Schnurr PP, et al. Cognitive behavioral therapy for posttraumatic stress disorder in women: a randomized controlled trial. *Journal of the American Medical Association*. 2007;297(8):820-30. DOI: 10.1001/jama.297.8.820. [Context Link 1] View abstract...
32. Garrison D, Daigler GE. Treatment settings for adolescent psychiatric conditions. *Adolescent Medicine Clinics* 2006;17(1):233-50. DOI: 10.1016/j.admed.2005.10.002. [Context Link 1, 2, 3] View abstract...
33. Timko C, Sempel JM, Moos RH. Models of standard and intensive outpatient care in substance abuse and psychiatric treatment. *Administration and Policy in Mental Health* 2003;30(5):417-36. [Context Link 1] View abstract...
34. Kallert TW, et al. Are day hospitals effective for acutely ill psychiatric patients? A European multicenter randomized controlled trial. *Journal of Clinical Psychiatry* 2007;68(2):276-87. [Context Link 1] View abstract...
35. Marshall M, Crowther R, Sledge WH, Rathbone J, Soares-Weiser K. Day hospital versus admission for acute psychiatric disorders. *Cochrane Database of Systematic Reviews* 2011, Issue 12. Art. No.: CD004026. DOI: 10.1002/14651858.CD004026.pub2. [Context Link 1] View abstract...
36. Stein DJ, Ipser JC, Seedat S. Pharmacotherapy for post traumatic stress disorder (PTSD). *Cochrane Database of Systematic Reviews* 2006, (verified by Cochrane 2009 Jan), Issue 1. Art. No.: CD002795. DOI: 10.1002/14651858.CD002795.pub2. [Context Link 1, 2, 3] View abstract...
37. Benedek DM, Friedman MJ, Zatzick D, Ursano RJ. Guideline watch: practice guideline for the treatment of patients with acute stress disorder and posttraumatic stress disorder. [Internet] American Psychiatric Association. 2009 Mar Accessed at: <http://psychiatryonline.org/guidelines.aspx>. (accessed 2012 Aug 28) DOI: 10.1176/appi.books.9780890423479. [Context Link 1, 2, 3]
38. Shalev AY. Posttraumatic stress disorder and stress-related disorders. *Psychiatric Clinics of North America* 2009;32(3):587-704. DOI: 10.1016/j.psc.2009.06.001. [Context Link 1, 2] View abstract...
39. Pae CU, et al. The atypical antipsychotics olanzapine and risperidone in the treatment of posttraumatic stress disorder: a meta-analysis of randomized, double-blind, placebo-controlled clinical trials. *International Clinical Psychopharmacology* 2008;23(1):1-8. DOI: 10.1097/YIC.0b013e32828aa324. [Context Link 1, 2, 3] View abstract...
40. Cohen JA, Mannarino AP, Peret JM, Staron V. A pilot randomized controlled trial of combined trauma-focused CBT and sertraline for childhood PTSD symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry* 2007;46(7):811-9. DOI: 10.1097/chi.0b013e3280547105. [Context Link 1, 2] View abstract...
41. Barrera TL, Mott JM, Hofstein RF, Teng EJ. A meta-analytic review of exposure in group cognitive behavioral therapy for posttraumatic stress disorder. *Clinical Psychology Review* 2012;33(1):24-32. DOI: 10.1016/j.cpr.2012.09.005. [Context Link 1] View abstract...
42. Bisson J, Andrew M. Psychological treatment of post-traumatic stress disorder (PTSD). *Cochrane Database of Systematic Reviews* 2007, (verified by Cochrane 2009 Jan), Issue 3. Art. No.: CD003388. DOI: 10.1002/14651858.CD003388.pub3. [Context Link 1, 2]
43. Ponniah K, Holton SD. Empirically supported psychological treatments for adult acute stress disorder and posttraumatic stress disorder: a review. *Depression and Anxiety* 2009;26(12):1086-109. DOI: 10.1002/da.20635. [Context Link 1, 2] View abstract...
44. Powers MB, Halpern JM, Forenchak MP, Gillham SJ, Foa EB. A meta-analytic review of prolonged exposure for posttraumatic stress disorder. *Clinical Psychology Review* 2010;30(6):635-41. DOI: 10.1016/j.cpr.2010.04.007. [Context Link 1, 2] View abstract...
45. Kowalik J, Weller J, Venter J, Drachman D. Cognitive behavioral therapy for the treatment of pediatric posttraumatic stress disorder: a review and meta-analysis. *Journal of Behavior Therapy and Experimental Psychiatry* 2011;42(3):405-13. DOI: 10.1016/j.jbtep.2011.02.002. [Context Link 1, 2] View abstract...
46. Macdonald G, et al. Cognitive-behavioural interventions for children who have been sexually abused. *Cochrane Database of Systematic Reviews* 2012, Art. No.: CD001930. DOI: 10.1002/14651858.CD001930.pub3. [Context Link 1, 2] View abstract...
47. Deblinger E, Mannarino AP, Cohen JA, Runyon MK, Steer RA. Trauma-focused cognitive behavioral therapy for children: impact of the trauma narrative and treatment length. *Depression and Anxiety* 2011;26(1):67-75. DOI: 10.1002/da.20744. [Context Link 1, 2] View abstract...
48. Practice guideline for the assessment and treatment of patients with suicidal behaviors. *American Journal of Psychiatry* 2003;160(11 Suppl):1-50. [Context Link 1, 2, 3]
49. Psychiatric evaluation of adults. Second edition. *American Journal of Psychiatry* 2006;163(6 Suppl):3-36. [Context Link 1, 2]
50. Zimmerman M, McGlinchey JB, Chelminski I, Young D. Diagnostic co-morbidity in 2300 psychiatric out-patients presenting for treatment evaluated with a semi-structured diagnostic interview. *Psychological Medicine* 2008;38(2):199-210. DOI: 10.1017/S0033291707001717. [Context Link 1, 2] View abstract...
51. Sowers WE, Rohland B. American Association of Community Psychiatrists' principles for managing transitions in behavioral health services. *Psychiatric Services* 2004;55(11):1271-5. [Context Link 1, 2] View abstract...

Footnotes

Return to top of *Posttraumatic Stress Disorder: Residential Care - BHG*

[A] Residential care is intended for patients who need around-the-clock behavioral care but do not need the high level of physical security and frequency of psychiatric and medical intervention that are available on an inpatient unit.(1)(2)(3) [A in Context Link 1, 2]

[B] Patients with persistent thoughts of suicide, homicide, or serious harm to self or another may need frequent monitoring for progression to planning or intention to act on their thoughts. Such monitoring can be safely conducted at an outpatient level of care if providers and supports are available and the patient can reliably participate in the monitoring process.(7) [B in Context Link 1, 2, 3, 4]

[C] In adults, randomized controlled trials have found that a variety of antidepressants from multiple medication classes are efficacious, with greatest efficacy demonstrated for SSRIs.(29)(36)(37)(38) A meta-analysis of studies examining the atypical antipsychotics olanzapine and risperidone in PTSD, either as monotherapy or as adjunctive pharmacotherapy, found them superior to placebo, although side effects, including weight gain, led to substantial dropout rates among those treated.(39) A randomized controlled trial of a pharmacotherapy for PTSD in children and adolescents compared sertraline and trauma-focused cognitive behavioral therapy (CBT) vs trauma-focused CBT alone and did not find that the combination therapy was more efficacious.(40) An expert consensus guideline recommends that an antidepressant be used to treat PTSD in a child or adolescent only if trauma-focused CBT has been unsuccessful.(29) [C in Context Link 1, 2, 3, 4]

[D] For PTSD in adults, meta-analyses of randomized controlled trials have found that individually administered as well as group trauma-focused cognitive behavioral therapy (CBT) and eye movement desensitization and reprocessing therapy were effective.(29)(41)(42)(43) A meta-analysis shows prolonged exposure therapy to be similarly effective.(44) A meta-analysis and systematic review of CBT for the treatment of PTSD in children

report that CBT was efficacious in this patient population.(45) A meta-analysis and systematic review of cognitive behavioral intervention for children who had been sexually abused reported that CBT had moderate efficacy in reducing PTSD and other anxiety symptoms.(46) A randomized trial comparing 8 weeks to 16 weeks of trauma-focused CBT in children found there was little extra improvement with 16 instead of 8 weeks of CBT.(47) [D in Context Link 1, 2, 3]

[E] See Clinical Indications for Admission to Residential Care in this guideline. [E in Context Link 1]

[F] Patient should be observed and safety checks should be performed as often as necessary to minimize risk of harm to self or others.(48) [F in Context Link 1]

[G] Precipitants explain why the admission occurred at the specific point in time. Exploration of all precipitants is essential for treatment and discharge planning.(49) [G in Context Link 1]

[H] PTSD has a high rate of comorbidity with other behavioral health disorders, particularly depressive disorders, substance use disorders, and other anxiety disorders.(30)(50) [H in Context Link 1, 2, 3]

[I] Psychosocial interventions should be individualized, address all admission precipitants and barriers to discharge, and involve family and other supports as necessary.(28) [I in Context Link 1, 2, 3]

[J] Pharmacotherapy should be accompanied by interactive psychoeducation that includes discussion of the course and treatment of PTSD and the potential beneficial and adverse effects of prescribed medication.(28)(29) [J in Context Link 1, 2]

[K] A plan for suicide, homicide, or serious harm to self or another involves more than just the identification of a method to be used, but also consideration of how the means of suicide, homicide, or harm will be acquired and where and when the action will be taken.(48) [K in Context Link 1]

[L] Harm to self or another is considered serious if it has a substantial likelihood of causing death, disability, or major disfigurement. [L in Context Link 1]

[M] A crisis plan establishes what actions the patient and supports are to take if dangerous ideation or behavior develops.(48) [M in Context Link 1, 2]

[N] Diagnostic tests may be performed off-site. [N in Context Link 1]

[O] On the basis of findings from meta-analyses of randomized controlled trials, a practice guideline recommends that an initial trial of trauma-focused cognitive behavioral therapy or eye movement desensitization and reprocessing for PTSD generally comprise 8 to 12 individual therapy sessions, each up to 90 minutes long, administered by the same therapist, on a weekly or more frequent basis. Hence, in some treatment settings a course of treatment may not be completed but may be planned or initiated.(29) [O in Context Link 1]

[P] Essential functions are those that are necessary to sustain life, such as feeding and hydrating oneself. [P in Context Link 1]

Codes

Return to top of Posttraumatic Stress Disorder: Residential Care - BHG

ICD-10 Diagnosis: F43.10, F43.11, F43.12

ICD-9 Diagnosis: 309.81

DSM-IV: 309.81

MCG™

Behavioral Health Care 17th Edition

Copyright © 2013 MCG Health, LLC

All Rights Reserved

Last Update: 1/26/2013 3:25:33 PM

Build Number: 17.0.47227.001287

Behavioral Health Care > Behavioral Health Level of Care Guidelines > Residential Acute Behavioral Health Level of Care, Child or Adolescent

Residential Acute Behavioral Health Level of Care, Child or Adolescent

ORG: B-802-RES (BHG)

MCGTM
Behavioral Health Care
17th Edition

- Admission Guidelines
- Recovery Course
- Discharge Guidelines
- Care Planning and Evaluation
- References
- Footnotes

Admission Guidelines

Return to top of Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG
(Expand All / Collapse All)

- Admission to Residential Acute Level of Care for Child or Adolescent is judged appropriate as indicated by **ALL** of the following (A):
 - Around-the-clock behavioral care is necessary for treatment because of **1 or more** of the following(4):
 - ☐ Imminent danger to self because of **1 or more** of the following(5)(6)(7):
 - Imminent risk for recurrence of Suicide attempt or act of serious self Harm as indicated by **ALL** of the following:
 - Very recent Suicide attempt or deliberate act of serious self Harm
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for suicide or serious self Harm
 - Command auditory hallucinations for suicide or serious self Harm
 - Dangerous behavior risk, persistent Thoughts of suicide or serious Harm to self, or suicide trigger state without formed thoughts, that cannot be adequately monitored at lower level of care because of **1 or more** of the following (5)(8):
 - Insufficient child or adolescent behavioral care (eg, facility, provider) availability
 - Severe conflict in family environment or other inadequacy in patient support system
 - Patient characteristics such as high impulsivity, unreliability, or extreme agitation with desperation
 - Ruminative flooding; uncontrollable and overwhelming profusion of negative thoughts
 - Frantic hopelessness; fatalistic conviction that life will not improve along with oppressive sense of entrapment and doom
 - ☐ Imminent danger to others due to **1 or more** of the following(9)(10)(11)(12):
 - Imminent risk for recurrence of an attempt to seriously Harm another as indicated by **ALL** of the following:
 - Very recent attempt to seriously Harm another
 - Absence of Sufficient relief of the action's precipitants
 - Current plan for homicide or serious Harm to another
 - Command auditory hallucinations or paranoid delusions contributing to risk for homicide or serious Harm to another
 - Persistent thoughts of, or violent impulsive acts that could likely result in, homicide or serious Harm to another that cannot be adequately monitored at lower level of care because of **1 or more** of the following (9):
 - Insufficient child or adolescent behavioral care (eg, facility, provider) availability
 - Severe conflict in family environment or other inadequacy in patient support system
 - Patient characteristics such as high impulsivity or unreliability
 - Life-threatening inability to receive adequate care from caretakers(9)(11)
 - Severe disability or disorder requiring acute residential intervention as indicated by **ALL** of the following:
 - Severe behavioral health disorder-related symptoms or condition are present as indicated by **1 or more** of the following(2)(3)(9)(13)(14)(15):
 - Major dysfunction in daily living (eg, family, interpersonal, school functioning)
 - Severe problem with cognition, memory, or judgment
 - Severe symptoms (eg, hallucinations, delusions, other acute psychotic symptoms, mania, severe autistic behaviors)(16)(17)(18)
 - Severe behavior risk (affective dysregulation) characteristics indicated by **1 or more** of the following:
 - Evidence of severely diminished ability to assess consequences of own actions (eg, acts of severe property damage)
 - Frequent extreme external (extreme angry outbursts) or internal (extreme sulking and rumination) anger manifestations
 - High levels of family conflict
 - Patient management at highest nonresidential level of care has failed or is not feasible until acute intervention or modification is initiated.
 - Severe comorbid substance use disorder that must be controlled (eg, abstinence necessary) to achieve stabilization of primary psychiatric disorder(19)(20)
 - Patient has currently stabilized during inpatient treatment stay for severe symptoms or behavior and requires structured setting with continued around-the-clock behavioral care.
 - No exclusions to treatment: situation and expectations are appropriate for level as indicated by **ALL** of the following(1)(2)(3)(4)(9)(5)(14)(21):

- Recommended treatment is necessary, appropriate, and not feasible at a lower level of care (ie, documented behavior, symptoms, or risk judged not appropriate for partial hospital, IOP, or acute outpatient care).
- Very short-term crisis intervention and resource planning for further care at a nonresidential level is unavailable or judged inappropriate.
- Patient has, at the least, minimal motivation to participate in treatment within a highly structured setting at the direction of parent or guardian.
- No anticipated need for physical restraint, seclusion, or other involuntary control (eg, patient not actively violent)
- No need for around-the-clock medical or nursing care
- Patient has sufficient cognitive capacity to respond to planned individual and group treatment components. (15)
- Adequate response (eg, stabilization for nonresidential level of care) to planned treatment is expected within a limited time period.

Recovery Course

Return to top of Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG

Stage	Clinical Status	Interventions	Evaluation
1	<ul style="list-style-type: none"> • Continued treatment needed for condition as described in Admission Guidelines • Treatment plan with goals and progress measurement in place • Response to treatment (progress toward goals) is likely 	<ul style="list-style-type: none"> • Appropriate crisis management instituted or documented as not needed • Ongoing care coordination • Ongoing discharge planning • Regular safety checks done at appropriate frequency • Appropriate treatment plan review 	<ul style="list-style-type: none"> • Evaluation completed and reviewed • Symptom and functioning assessment at appropriate frequency documented • Physician evaluation of patient and progress at appropriate frequency documented • Review of need for medication or adjustment at appropriate frequency documented
2	<ul style="list-style-type: none"> • Risk status acceptable • Functional status acceptable 	<ul style="list-style-type: none"> • Medical needs manageable 	

(2)(3)(4)(5)(15)(22)(23)

Recovery Milestones are indicated in bold.

Discharge Guidelines

Return to top of Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG
[Expand All / Collapse All]

- Continued residential care is generally needed until **1** or more of the following(21):
 - Residential care no longer necessary due to adequate patient stabilization or improvement as indicated by **ALL** of the following(4)(5)(22):
 - ☐ Risk status acceptable as indicated by **ALL** of the following:
 - Patient has not recently made a Suicide attempt or act of serious Harm to self, or has had Sufficient relief of precipitants of any such action.
 - Absence of Current plan for suicide or serious Harm to self for at least 24 hours
 - Thoughts of suicide, homicide, or serious Harm to self or to another are absent or manageable at available lower level of care.
 - Supports, and patient as appropriate, understand follow-up treatment and crisis plan.
 - Provider and supports are sufficiently available at lower level of care.
 - Patient, as appropriate, can participate as needed in monitoring at next level of care.
 - ☐ Functional status acceptable as indicated by **1** or more of the following:
 - No essential function (9) is significantly impaired.
 - An essential function is impaired, but impairment is manageable at available lower level of care.
 - ☐ Medical needs manageable as indicated by **ALL** of the following:
 - Adverse medication effects absent or manageable at available lower level of care
 - Medical comorbidity absent or manageable at available lower level of care
 - Substance withdrawal absent or manageable at available lower level of care
 - Residential care no longer appropriate due to patient progress record or consent as indicated by **1** or more of the following:
 - Patient deterioration requires higher level of care.
 - Guardian no longer consents to treatment.

Care Planning and Evaluation

Return to top of Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG

- Planning and evaluating appropriate care should address:
 - Evaluation and care needs, which may include(24)(25)(26)(27):
 - Treatments and procedures, including:
 - Psychosocial interventions emphasizing admission precipitants and barriers to discharge

- Cognitive behavioral therapy, interpersonal psychotherapy, or other evidence-based psychosocial therapy appropriate for diagnosis
- Parental component of psychosocial interventions
- Clinical management and psychoeducation if medication is prescribed
- Psychotropic medication if indicated
- Consultation, assessment, and other services, including:
 - Substance abuse assessment(24)
 - Social services consultation for placement or housing
 - Conference with school
- Discharge planning needs, which may include(28):
 - Rapid planning for next level of care; considerations include:
 - Plan for monitoring for dangerous ideation or behavior if necessary
 - Plan for assisting patient with self-care if necessary
 - Aftercare plan development with input from multiple providers, patient, and patient's supports
 - Preparation of patient and supports for transition to next level of care, including:
 - Ensure sufficient knowledge of:
 - Patient's illness
 - Medication
 - Risk factors for relapse
 - Warning signs of relapse
 - Assess ability to attend school, work (as appropriate), and participate in usual activities.
 - Review crisis plan with patient and supports.
 - Follow-up appointments, including:
 - Psychiatrist for pharmacotherapy and clinical management
 - Therapist for cognitive behavioral therapy, interpersonal psychotherapy, or other evidence-based psychosocial therapy appropriate for diagnosis
 - Medical care visit (eg, primary care)
 - Referrals for community assistance and support, including:
 - Self-help or support groups for patient, family, and caregivers
 - Community services for housing, financial, or transportation needs
 - Medications and supplies, including:
 - Psychotropic medications
 - Medications for comorbid medical conditions

References

Return to top of *Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG*

1. Lloyd-Evans B, Slade M, Jagielska D, Johnson S. Residential alternatives to acute psychiatric hospital admission: systematic review. *British Journal of Psychiatry* 2009;195(2):109-17. DOI: 10.1192/bjp.bp.108.058347. [Context Link 1, 2] View abstract...
2. Mee-Lee D, Shulman GD, Fishman M, Gastfriend DR, Griffith JH. ASAM Patient Placement Criteria for the Treatment of Substance-Related Disorders, Second Edition-Revised, 2nd rev ed. Chevy Chase, MD: American Society of Addiction Medicine; 2001. [Context Link 1, 2, 3, 4]
3. Kleber HD, et al. Practice guideline for the treatment of patients with substance use disorders. 2nd ed. [Internet] American Psychiatric Association; 2007 Apr. Accessed at: <http://psychiatryonline.org/guidelines.aspx>. [created 2006 May; accessed 2012 Sep 20] DOI: 10.1176/appi.books.9780890423363.141077. [Context Link 1, 2, 3, 4]
4. Kober D, Martin A. Inpatient psychiatric, partial hospital, and residential treatment for children and adolescents. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3766-71. [Context Link 1, 2, 3, 4]
5. Wagner KD, Brent DA. Depressive disorders and suicide. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3652-62. [Context Link 1, 2, 3, 4, 5, 6]
6. Screening and treatment for major depressive disorder in children and adolescents: US Preventive Services Task Force Recommendation Statement. *Pediatrics* 2009;123(4):1223-8. DOI: 10.1542/peds.2008-2381. [Context Link 1, 2] View abstract...
7. Olesvold T, Bakkejord T, Hansen V, Nivison M, Sorgaard KW. Suicidality related to first-time admissions to psychiatric hospital. *Social Psychiatry and Psychiatric Epidemiology* 2012;47(3):419-25. DOI: 10.1007/s00127-011-0343-2. [Context Link 1] View abstract...
8. Yaseen Z, Katz C, Johnson MS, Eisenberg D, Cohen LJ, Galynker II. Construct development: The Suicide Trigger Scale (STS-2), a measure of a hypothesized suicide trigger state. *BMC Psychiatry* 2010;10:110. DOI: 10.1186/1471-244X-10-110. [Context Link 1] View abstract...
9. Lichtenberg P. The residential care alternative for the acutely psychotic patient. *Psychiatric Quarterly* 2011;82(4):329-41. DOI: 10.1007/s11126-011-9176-0. [Context Link 1, 2, 3, 4] View abstract...
10. Meuteners LB, Lee AH, Hendrie D. Interpersonal violence hospitalisations for adolescents: a population-based study. *Journal of Paediatrics and Child Health* 2010;46(11):886-90. DOI: 10.1111/j.1440-1754.2010.01818.x. [Context Link 1] View abstract...
11. Baron DA, Dubin WR, Ning A. Other psychiatric emergencies. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:2732-45. [Context Link 1, 2]
12. Myklebust LH, Sorgaard K, Rotvold K, Wynn R. Factors of importance to involuntary admission. *Nordic Journal of Psychiatry* 2012;66(3):176-82. DOI: 10.3109/08039468.2011.611252. [Context Link 1] View abstract...
13. Brooker C, Ricketts T, Bennett S, Lemme F. Admission decisions following contact with an emergency mental health assessment and intervention service. *Journal of Clinical Nursing* 2007;16(7):1313-22. DOI: 10.1111/j.1365-2702.2007.01302.x. [Context Link 1] View abstract...
14. Stewart SE, et al. Long-term outcome following intensive Residential Treatment of Obsessive-Compulsive Disorder. *Journal of Psychiatric Research* 2009;43(13):1118-23. DOI: 10.1016/j.jpsychires.2009.03.012. [Context Link 1, 2] View abstract...
15. Boyer DE, Kane C. Program evaluation of a community crisis stabilization program. *Archives of Psychiatric Nursing* 2010;24(6):387-96. DOI: 10.1016/j.apnu.2010.03.002. [Context Link 1, 2, 3] View abstract...

16. Fochtmann LJ, Mojtabai R, Bromet EJ. Other psychotic disorders. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:1605-28. [Context Link 1]
17. Volkmar FR, Klin A, Schultz RT, Stale MW. Pervasive developmental disorders. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:3540-59. [Context Link 1]
18. Lewis S, Escalona PR, Keith SJ. Phenomenology of schizophrenia. In: Sadock BJ, Sadock VA, Ruiz P, editors. *Kaplan & Sadock's Comprehensive Textbook of Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:1433-50. [Context Link 1]
19. Kuehn BM. Integrated care key for patients with both addiction and mental illness. *Journal of the American Medical Association* 2010;303(19):1905-7. DOI: 10.1001/jama.2010.597. [Context Link 1]
20. Torrens M, Gilchrist G, Domingo-Salvany A, psyCoBarcelona Group. Psychiatric comorbidity in illicit drug users: substance-induced versus independent disorders. *Drug and Alcohol Dependence* 2011;113(2-3):147-56. DOI: 10.1016/j.drugalcdep.2010.07.013. [Context Link 1] View abstract...
21. Hawthorne WB, et al. A randomized trial of short-term acute residential treatment for veterans. *Psychiatric Services* 2005;56(11):1379-86. DOI: 10.1176/appi.ps.56.11.1379. [Context Link 1, 2] View abstract...
22. Sharfstein SS. Goals of inpatient treatment for psychiatric disorders. *Annual Review of Medicine* 2009;60:393-403. DOI: 10.1146/annurev.med.60.042607.080257. [Context Link 1, 2] View abstract...
23. Bellonci C. Physician leadership in residential treatment for children and adolescents. *Child and Adolescent Psychiatric Clinics of North America* 2010;19(1):21-30; table of contents. DOI: 10.1016/j.chc.2009.08.001. [Context Link 1] View abstract...
24. Psychiatric evaluation of adults. Second edition. *American Journal of Psychiatry* 2006;163(6 Suppl):3-36. [Context Link 1, 2]
25. Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. *Cochrane Database of Systematic Reviews* 2011, Issue 9. Art. No.: CD005031. DOI: 10.1002/14651858.CD005031.pub4. [Context Link 1] View abstract...
26. Soyka M, et al. The World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the biological treatment of substance use and related disorders. Part 2: Opioid dependence. *World Journal of Biological Psychiatry* 2011;12(3):160-87. DOI: 10.3109/15622975.2011.561872. [Context Link 1] View abstract...
27. Winters KC, Botzet AM, Fahnhorst T. Advances in adolescent substance abuse treatment. *Current Psychiatry Reports* 2011;13(5):416-21. DOI: 10.1007/s11920-011-0214-2. [Context Link 1] View abstract...
28. Sowers WE, Rohland B. American Association of Community Psychiatrists' principles for managing transitions in behavioral health services. *Psychiatric Services* 2004;55(11):1271-5. [Context Link 1] View abstract...

Footnotes

Return to top of *Residential Acute Behavioral Health Level of Care, Child or Adolescent - BHG*

[A] Residential care is intended for patients who need around-the-clock behavioral care but do not need the high level of physical security and frequency of psychiatric and medical intervention that are available on an inpatient unit.(1)(2)(3) [A in Context Link 1]

[B] Patients with persistent thoughts of suicide, homicide, or serious harm to self or another may need frequent monitoring for progression to planning or intention to act on their thoughts. Such monitoring can be safely conducted at an outpatient level of care if providers and supports are available and the patient can reliably participate in the monitoring process.(5)(6) [B in Context Link 1, 2]

[C] Essential functions are those that are necessary to sustain life, such as feeding and hydrating oneself. [C in Context Link 1]

MCG™
Behavioral Health Care 17th Edition
Copyright © 2013 MCG Health, LLC
All Rights Reserved

Last Update: 1/26/2013 3:27:29 PM
Build Number: 17.0.47227.001287